Phone: 415-863-2380 Support: info@quickarrays.com Web: www.quickarrays.com

Advancing Biomedical Science Through Tissue Amara.

Cat No: **THC481** - Thyroid cancer tissue array with paired normal tissues, 48 samples (2mm)

| Lot# | Cores | Size | Cut | Format | QA/QC | | 12345678 |
|----------|-------|------|-----|--------|-------------------------------|-----------------|-----------------------------------------|
| THC48101 | 48 | 2mm | 4um | 6X8 | H&E, IHC anti- Cytokeratin | QUICKARRAYS, IN | A S S S S S S S S S S S S S S S S S S S |

Recommended applications: For research use only. Designed for thyroid normal vs cancer RNA or protein tissue profiling using IHC or ISH.

Description: Thyroid cancer tissue array, 16 cases, 48 cores, one normal paired with two tumor tissue cores from each patient with grading and TNM staging data.

All the tissues were from surgical resection. They were fixed in 10% neutral buffered formalin for 24 hours and processed using identical SOPs. Sections were picked onto Superfrost Plus or APES coated Superfrost slides. They can be stored for use at 4C for up to six months from the date of shipment. There may be 5 to 10% of tissue core loss.

| Array Position | Sex | Age | Anatomic Site | Pathology | Grade | Stage |
|----------------|-----|-----|---------------|-------------------------------------|-------|--------|
| A01 | F | 64 | Thyroid | Papillary adenocarcinoma | null | T2N1M0 |
| A02 | М | 51 | Thyroid | Papillary adenocarcinoma | null | T1N0M0 |
| A03 | М | 58 | Thyroid | Papillary adenocarcinoma | null | T1N0M0 |
| A04 | F | 30 | Thyroid | Papillary adenocarcinoma | null | T2N0M0 |
| A05 | М | 37 | Thyroid | Papillary adenocarcinoma | null | T2N0M0 |
| A06 | F | 56 | Thyroid | Papillary adenocarcinoma | null | T1N0M0 |
| A07 | F | 66 | Thyroid | Follicular carcinoma | null | T2N0M0 |
| A08 | F | 16 | Thyroid | Papillary adenocarcinoma | null | T3N1M0 |
| B01 | F | 64 | Thyroid | Papillary adenocarcinoma | null | T2N1M0 |
| B02 | М | 51 | Thyroid | Papillary adenocarcinoma | null | T1N0M0 |
| B03 | М | 58 | Thyroid | Papillary adenocarcinoma | null | T1N0M0 |
| B04 | F | 30 | Thyroid | Papillary adenocarcinoma | null | T2N0M0 |
| B05 | М | 37 | Thyroid | Papillary adenocarcinoma | null | T2N0M0 |
| B06 | F | 56 | Thyroid | Papillary adenocarcinoma | null | T1N0M0 |
| B07 | F | 66 | Thyroid | Follicular carcinoma | null | T2N0M0 |
| B08 | F | 16 | Thyroid | Papillary adenocarcinoma | null | T3N1M0 |
| C01 | F | 64 | Thyroid | Uninvolved thyroid tissue of A1, B1 | null | null |
| C02 | М | 51 | Thyroid | Uninvolved thyroid tissue of A2, B2 | null | null |
| C03 | М | 58 | Thyroid | Uninvolved thyroid tissue of A3, B3 | null | null |
| C04 | F | 30 | Thyroid | Uninvolved thyroid tissue of A4, B4 | null | null |
| C05 | М | 37 | Thyroid | Uninvolved thyroid tissue of A5, B5 | null | null |
| C06 | F | 56 | Thyroid | Uninvolved thyroid tissue of A6, B6 | null | null |
| C07 | F | 66 | Thyroid | Uninvolved thyroid tissue of A7, B7 | null | null |
| C08 | F | 16 | Thyroid | Uninvolved thyroid tissue of A8, B8 | null | null |
| D01 | F | 27 | Thyroid | Papillary adenocarcinoma | null | T3N1M0 |
| D02 | F | 49 | Thyroid | Papillary adenocarcinoma | null | T4N0M0 |
| D03 | F | 27 | Thyroid | Papillary adenocarcinoma | null | T2N0M0 |
| D04 | F | 19 | Thyroid | Follicular papillary adenocarcinoma | null | T2N1M0 |
| D05 | М | 34 | Thyroid | Follicular papillary adenocarcinoma | null | T2N1M0 |
| D06 | М | 16 | Thyroid | Follicular carcinoma | null | T3N1M0 |
| D07 | F | 35 | Thyroid | Follicular carcinoma | null | T3N0M0 |
| D08 | М | 60 | Thyroid | Follicular carcinoma | null | T3N0M0 |
| E01 | F | 27 | Thyroid | Papillary adenocarcinoma | null | T3N1M0 |
| E02 | F | 49 | Thyroid | Papillary adenocarcinoma | null | T4N0M0 |
| E03 | F | 27 | Thyroid | Papillary adenocarcinoma | null | T2N0M0 |
| E04 | F | 19 | Thyroid | Follicular papillary adenocarcinoma | null | T2N1M0 |
| E05 | М | 34 | Thyroid | Follicular papillary adenocarcinoma | null | T2N1M0 |
| E06 | М | 16 | Thyroid | Follicular carcinoma | null | T3N1M0 |
| E07 | F | 35 | Thyroid | Follicular carcinoma | null | T3N0M0 |
| E08 | М | 60 | Thyroid | Follicular carcinoma | null | T3N0M0 |

Quickarrays Inc., 1049 Union Ave, Suite F, Fairfield, CA 94533 Phone: 415-863-2380; Fax: 510-225-1084; Email: info@quickarrays.com

Phone: 415-863-2380 Support: info@quickarrays.com Web: www.quickarrays.com

| | Advancing Biomedical Science Through Tissue Arraya | | | | | | |
|-----|----------------------------------------------------|----|---------|-------------------------------------|------|------|--|
| F01 | F | 27 | Thyroid | Uninvolved thyroid tissue of D1, E1 | null | null | |
| F02 | F | 49 | Thyroid | Uninvolved thyroid tissue of D2, E2 | null | null | |
| F03 | F | 27 | Thyroid | Uninvolved thyroid tissue of D3, E3 | null | null | |
| F04 | F | 19 | Thyroid | Uninvolved thyroid tissue of D4, E4 | null | null | |
| F05 | М | 34 | Thyroid | Uninvolved thyroid tissue of D5, E5 | null | null | |
| F06 | М | 16 | Thyroid | Uninvolved thyroid tissue of D6, E6 | null | null | |
| F07 | F | 35 | Thyroid | Uninvolved thyroid tissue of D7, E7 | null | null | |
| F08 | М | 60 | Thyroid | Uninvolved thyroid tissue of D8, E8 | null | null | |

Notes: Bake at 60C for ~60 minutes before use. If antigen retrieving is needed, it is important to avoid **direct-boiling and high pH or high strength** antigen retrieving buffer.

Certified by: Langxing Pan, M.D.

TNM Classification: Thyroid carcinoma

T- Primary tumor

- TX Primary tumor cannot be assessed
- TO No evidence of primary tumor
- T1 Tumor 1 cm or less in greatest dimension, limited to the thyroid
- T2 Tumor more than 1 cm but not more than 4 cm in greatest dimension, limited to the thyroid
- T3 Tumor more than 4 cm in greatest dimension, limited to the thyroid
- T4 Tumor of any size extending beyond the thyroid capsule

N - Regional lymph nodes

- NX Regional lymph nodes cannot be assessed
- NO No regional lymph node metastasis
- N1 Regional lymph node metastasis
 - N1a Metastasis in ipsilateral cervical lymph nodes
 - N1b Metastasis in bilateral, midline, or contralateral cervical or mediastinal lymph nodes

M - Distant metastasis

- MX Distant metastasis cannot be assessed
- MO No distant metastasis
- M1 Distant metastasis

Histopathologic Types

- Papillary carcinoma
- Follicular carcinoma
- Medullary carcinoma
- Undifferentiated (anaplastic) carcinoma

TNM classification of malignant tumours, Fifth Edition (1997)